

## ***In vitro* Ubiquitin Dimer Formation Assay**

Sheng Wang<sup>1</sup>, Ling Cao<sup>1,2</sup> and Hong Wang<sup>1,\*</sup>

<sup>1</sup>Department of Biochemistry, University of Saskatchewan, Saskatoon, Canada; <sup>2</sup>National Key laboratory of Crop Genetic Improvement, College of Plant Science and Technology, Huazhong Agricultural University, Wuhan, China

\*For correspondence: [hong.wang@usask.ca](mailto:hong.wang@usask.ca)

**[Abstract]** The process of protein ubiquitination typically consists of three sequential steps to add an ubiquitin (Ub) or Ub chain to a substrate protein, requiring three different enzymes, ubiquitin activating enzyme (E1), ubiquitin conjugating enzyme (E2), and ubiquitin protein ligase (E3). Most E2s possess the classical E2 activity in forming E2-Ub complex through a thioester linkage, in presence of an E1 and Ub. Additionally, some E2s have the ability of catalyzing the formation of free Ub dimer. Such activity indicates an important role of these E2s in ubiquitination pathway. Thus, we developed an *in vitro* Ub dimer formation assay to determine the activity of certain E2s. Moreover, by using Ub mutants, in which different lysine residues are mutated, the specific linkage of dimer can also be determined.

**Keywords:** Ubiquitination, Ubiquitin dimer formation, E2, *Arabidopsis* UBC22, K11 linkage

**[Background]** The existing protocols for E2 conjugation initiation assay (without adding E3 and substrate) aim to detect the thioester linkage (E2-S-Ub). Our method focuses on the E2 activity of catalyzing free Ub dimer formation (Ub-Ub). It provides a convenient way to detect an important biochemical feature of E2 in different species. Further, the specific linkage of dimer can be determined by using different Ub mutants.

### **Materials and Reagents**

1. 1.5 ml polypropylene tubes
2. Polyvinylidene difluoride (PVDF) membrane (Bio-Rad Laboratories, catalog number: 162-0177)
3. AmershamHyperfilm™ ECL (GE Healthcare, catalog number: 28906836)
4. Purified human recombinant E1 (BOSTONBIOCHEM, catalog number: K-995)
5. Qiagen Ni-NTA Spin Kit (QIAGEN, catalog number: 31314)
6. Purified human recombinant Ub (BOSTONBIOCHEM, catalog number: K-995)
7. Purified human recombinant Ub with the lysine 11 (K11) residue mutated (Ub-K11R) (BOSTONBIOCHEM, catalog number: UM-K11R)
8. Purified human recombinant Ub with the lysine 48 (K48) residue mutated (Ub-K48R) (BOSTONBIOCHEM, catalog number: UM-K48R)
9. Purified human recombinant Ub with the lysine 63 (K63) residue mutated (Ub-K63R) (BOSTONBIOCHEM, catalog number: UM-K63R)

10. 10x reaction buffer (BOSTONBIOCHEM, catalog number: K-995)
11. Mg-ATP solution (BOSTONBIOCHEM, catalog number: K-995)
12. 4x non-reducing loading buffer (BOSTONBIOCHEM, catalog number: K-995)
13. SDS-PAGE gel
14. Skimmed milk powder
15. Anti-ubiquitin antibody (Cell Signaling Technology, catalog number: 3936)
16. Goat anti-mouse antibody conjugated to horseradish peroxidase (HRP) (Bio-Rad Laboratories, catalog number: 170-6516)
17. Amersham ECL prime Western blotting detection reagent (GE Healthcare, catalog number: RPN2232)
18. NaCl
19. KCl
20. Na<sub>2</sub>HPO<sub>4</sub>
21. KH<sub>2</sub>PO<sub>4</sub>
22. Tween-20 (Sigma-Aldrich, catalog number: P1379)
23. Tris base
24. Glycine
25. Methanol
26. Dialysis buffer (see Recipes)
27. 1x PBS (see Recipes)
28. 1x PBST (see Recipes)
29. Transfer buffer (see Recipes)

## **Equipment**


1. Incubator (VWR, model number: 1545) or water bath
2. Protein electrophoresis apparatus (Bio-Rad Laboratories, model: Mini PROTEAN® 3 Cell)
3. Western blotting apparatus (Bio-Rad Laboratories, model: Mini Trans-Blot® Cell)
4. X-Ray film processor (PROTEC, model: OPTIMAX)

## **Procedure**

1. Purify 6x His-tagged *Arabidopsis* E2 UBC22 (His-UBC22) using QIAGEN Ni-NTA Spin Kit and following the manufacturer's instructions. Purified protein is then dialyzed in a dialysis buffer for overnight.
2. Prepare the reaction samples in 1.5 ml polypropylene tubes. Different components are added as listed below. To test specific linkage of dimer, wild-type Ub can be substituted by different Ub mutants, each of which has one particular lysine residue mutated (Ub-K11R, Ub-K48R or Ub-K63R).

Component	Amount to add or final concentration
10x buffer	2.0 $\mu$ l
E1	0.2 $\mu$ M
E2	5.0 $\mu$ M
Ub	62.5 $\mu$ M
Mg-ATP solution	1.0 mM
ddH <sub>2</sub> O	make up to 20 $\mu$ l

3. Incubate the reaction samples at 30 °C either in a water bath or an incubator for 4 h.
4. Add 7  $\mu$ l 4x SDS non-reducing loading buffer and 1  $\mu$ l ddH<sub>2</sub>O into each reaction sample.
5. Treat the samples at 95 °C for 5 min.
6. Bio-Rad Mini PROTEAN® 3 Cell gel apparatus is used for protein electrophoresis. Load half amount of samples on a 15% SDS-PAGE resolving gel with 4% stacking gel and run the gel at 120 V for 130 min to separate the proteins.
7. Transfer the proteins to a piece of PVDF membrane at 120 V for 120 min using Bio-Rad Mini Trans-Blot® Cell. The transfer apparatus is placed into a bucket with ice to reduce heat production during the transfer.
8. Incubate the membrane in 10 ml of 1x PBS containing 5% skimmed milk powder for 0.5 h at room temperature for the blocking treatment.
9. Incubate the membrane with primary antibody (anti-ubiquitin antibody, antibody dilution ratio: 1:40,000) in 10 ml 1x PBST containing 3% skimmed milk powder overnight at 4 °C.
10. Wash the membrane with 1x PBST for three times at room temperature (20 ml, 10 min each).
11. Incubate the membrane with secondary antibody (goat anti-mouse antibody conjugated to HRP, antibody dilution ratio: 1:7,000) in 1x PBST containing 3% skimmed milk powder for 1 h at room temperature.
12. Wash the membrane with 1x PBST for three times at room temperature (20 ml, 10 min each).
13. Detect signal with the ECL prime Western blotting detection reagent according to the manufacturer's instructions (Figure 1). Use 1-10 min as the initial range of film exposure time, depending on signal strength.



**Figure 1. Ub dimer formation assay by His-UBC22.** To determine the activity of *Arabidopsis* UBC22, various components, as indicated on the top of the figure, were added to the reaction tubes with (lanes 2, 4, 6 and 8) or without His-UBC22 (lanes 1, 3, 5 and 7). The reaction tubes were incubated at 30 °C for 4 h. The samples were then subjected to SDS-PAGE (with 4% stacking and 15% resolving gel). Free Ub and Ub dimers were detected by Western blotting using an anti-Ub antibody. Free Ub (Ub) and Ub dimers (Ub<sub>2</sub>) are indicated on the right side of the figure. For the K11R mutant, no new Ub dimer formation was observed when His-UBC22 protein was added. Due to slight impurity, there was one weak band in the recombinant K11R protein which is slightly higher than the Ub dimer synthesized from the wild-type Ub and other Ub mutants. Three independent experiments were performed and had similar results (Figure from Wang *et al.*, 2016; Figure 6 in the manuscript: <http://jxb.oxfordjournals.org/content/early/2016/04/10/jxb.erw142.full>).

### Data analysis

Purified *Arabidopsis* recombinant E2 UBC22 fused with 6x His tag (His-UBC22) was tested in the *in vitro* Ub dimer formation assay. As shown in Figure 1, Ub dimers could be detected when His-UBC22 was added into the reaction (lane 2 compared to the lane 1), indicating the biochemical activity of UBC22 catalyzing free dimer formation. In addition, similar dimer formation was observed when the Ub-K48R mutant or Ub-K63R mutant was used, which lacks K48 residue or K63 residue (lane 6 compared to lane 5, or lane 8 compared to lane 7). Interestingly, when the Ub-K11R which lacks K11 residue was used, little dimer was produced (lane 4 compared to lane 3). These results indicate that UBC22 could catalyze Ub dimer formation *in vitro* specifically through K11 residue of Ub. Three independent experiments were performed and produced similar results.

## **Notes**

1. This assay aims to investigate a specific activity of the E2 enzyme. The purity of the recombinant protein or any conditions affecting a protein's activity could affect the result. A commercial E2 protein, such as Human E2 Ube2S, may be good for a positive control.
2. Poly-Ub chains might be formed and detected in the assay depending on the activity of an E2 tested.

## **Recipes**

1. Dialysis buffer  
10 mM Tris-HCl (pH 7.5)  
50 mM NaCl
2. 1x PBS  
137 mM NaCl  
2.7 mM KCl  
10 mM Na<sub>2</sub>HPO<sub>4</sub>  
1.8 mM KH<sub>2</sub>PO<sub>4</sub>  
Adjust to pH 7.4
3. 1x PBST  
0.1% Tween-20 in 1x PBS
4. Transfer buffer  
25 mM Tris base  
192 mM glycine  
20% (v/v) methanol

## **Acknowledgments**

We gratefully acknowledge the financial support from the Natural Sciences and Engineering Research Council of Canada (NSERC) (Discovery grant) to HW.

## **Reference**

1. Wang, S., Cao, L. and Wang, H. (2016). [Arabidopsis ubiquitin-conjugating enzyme UBC22 is required for female gametophyte development and likely involved in Lys11-linked ubiquitination.](#) *J Exp Bot* 67(11): 3277-3288.