

## Trichome Isolation and Integrity Test from *Brassica villosa* and Other Species

Naghabushana Nayidu<sup>1</sup>, Peta Bonham-Smith<sup>1</sup> and Margaret Y. Gruber<sup>2\*</sup>

<sup>1</sup>Biology Department, University of Saskatchewan, Saskatoon, Canada; <sup>2</sup>Agriculture and Agri-Food Canada, Saskatoon, Canada

\*For correspondence: [nagabushana@gmail.com](mailto:nagabushana@gmail.com); [Margie.Gruber@agr.gc.ca](mailto:Margie.Gruber@agr.gc.ca)

**[Abstract]** The outward growths from one or more epidermal cells are well known as trichomes (plant hair cells) (Levin, 1973; Mathur, 2006). Preparation of pure intact non-glandular trichomes from trichome-rich *Brassica villosa* depended on rendering the trichomes sufficiently stiff to dislodge them from the leaves in an undamaged state, which can be further used for structural, transcriptome, genome and biochemical or chemical analysis. Dislodging the flexible trichomes from *Brassica villosa* (*B. villosa*) with a paintbrush [as in Zhang and Oppenheimer (2004)] proved too gentle, and scraping trichomes off the leaves with a razor blade [as in Zhang and Oppenheimer (2004)] resulted in trichome cell disruption. Aziz *et al.* (2005) reported isolating alfalfa glandular trichomes by shearing in liquid nitrogen (N<sub>2</sub>). In the present study, a similar method was used. Non-glandular leaf trichomes were isolated by treating tissue with liquid N<sub>2</sub> to stiffen the flexible trichomes, followed by 1 min of shearing force with a common vortex mixer to dislodge the trichomes from their leaf bed (Nayidu *et al.*, 2014). Up-to ~20 % of the trichomes were removed from the leaf surface and the majority were unbroken (confirmed by no staining by trypan blue; Figure 1). Trichomes were then purified by straining a tissue/trichome/water mixture through a sieve. However, leaf tissues also became very brittle after dipping in liquid N<sub>2</sub> and broke up into very small pieces if the leaf tissue was agitated for a longer time period. Hence, N<sub>2</sub>-treated leaf samples could not be re-used to recover remaining attached trichomes. Contaminating leaf pieces from a 1 min shear were larger and easily to separate manually from the detached trichomes within the sieve, rendering a purified intact trichome preparation. The method was also successful at purifying trichomes from soybean (*Glycine max*) and tomato (*Solanum lycopersicum*).

### **Materials and Reagents**

1. Fresh trichome-bearing leaf tissue from *Brassica villosa* (drepansis cv.) (from two month old up-to two year leaves), soybean cv. B220, and tomato cv. Roma

*Note: Depending on the plant species and the subsequent type of analysis after trichome isolation, any tissue with non-glandular trichomes will likely suffice. However, tissues with denser trichome coverage will provide a greater yield of detached trichomes.*

2. Liquid nitrogen
- Note: Training in the safe use of liquid nitrogen is imperative.*
3. Sterile Millipore purified water
4. Ice
5. 0.4 % trypan blue (see Recipes)

### **Equipment**

1. Gloves to resist freezing temperatures (Cryo-Gloves) (VWR International, catalog number: 32885-736)
2. Polyfoam insulated box with ice
3. 1-ml pipette with plastic pipette tips
4. 50 ml BD Falcon™ conical plastic centrifuge tubes (BD, catalog number: 14-432-22) or plastic 250 ml centrifuge bottles (Thermo Fisher Scientific, catalog number: EW-06105-20)
5. BD-Falcon nylon mesh cell strainer (40  $\mu$ m pore size) (BD Biosciences, catalog number: 08-771-1)
6. Blunt end forceps to remove contaminating debris from the strainer (Seton, catalog number: AA969)
7. Genie 2 vortex mixer (Thermo Fisher Scientific) having round mixer head (for tubes) and flat mixer head (for larger bottles)
8. -80 °C freezer
9. 4 °C refrigerator
10. Flexidry™ unit (FTS Systems, catalog number: 4283)
11. Controlled environment greenhouse or growth cabinet.

*Note: Plant growth conditions depend on the nature of subsequent analyses. For example, chemical composition of plant tissues can change substantially with altered light intensity and temperature (16 h light/8 h dark, at least 400  $\mu$ E.m<sup>-2</sup>.s<sup>-1</sup>, 20/17 °C recommended, but light intensity from 700-1,400  $\mu$ E.m<sup>-2</sup>.s<sup>-1</sup> will mimic outside daylight).*

12. Axiovert 100 microscope (ZEISS) with 10x magnification

### **Procedure**

1. Healthy trichome-bearing leaves of *B. villosa*, soybean, and tomato were selected after growth in a greenhouse.

*Note: For *B. villosa*, up to 2 year old plant leaves can be used.*

2. Harvested leaves were cut into different sizes based on the size of the vessel used for the isolation of trichome (see Video 1 below).

*Note: One 3 cm x 3 cm leaf piece if using a 50 ml tube and two 10 cm x 4 cm strips if using 250 ml centrifuge bottles. The choice of tube and tissue size is dependent on the amount of trichome tissue needed for subsequent analyses (see recommendations below). For any leaf size used, the maximum purity trichome preparation achieved is ~20% of the total trichomes present on the leaf piece.*

3. Maximum 30% of the tube or bottle was filled with liquid N<sub>2</sub> and capped (loosely screwed on) to prevent overflow during mixing (without tube breakage from N<sub>2</sub> vapour pressure).
4. Using gloves, tubes were mixed for 1 min in liquid N<sub>2</sub> using a Genie 2 vortex mixer at maximum speed (level 8). One min vortexing maximizes the recovery of detached trichomes while minimizing the disruption of leaf tissue into small contaminating fragments that are difficult to remove later.
5. The tubes were then kept fully open on ice to maximize liquid N<sub>2</sub> evaporation.
6. Large leaf pieces were manually removed from the tubes using forceps and retained as a trichome-stripped leaf control sample.
7. Released trichomes adhering to the inside walls of the tube were gently suspended by swirling in 2 ml of sterile Millipore purified water.
8. Water initially frozen in the tube (because of liquid N<sub>2</sub>) was kept on ice for 3-4 min to thaw.
9. Thawed water from the tubes having trichomes (and also minute pieces of leaves) was sieved through a BD-Falcon nylon mesh cell strainer into a fresh 50 ml tubes.
10. Minute pieces of leaves on the cell strainer netting were again manually removed using forceps.
11. The strainer was then inverted over the same fresh tube.
12. Trichomes were dislodged into the tube by gently sweeping them (i.e. rinsing the strainer with a pipette) using an additional 1-2 ml of Millipore water (see video 1 below).

*Note: Cell strainers could be used up to 8-10 times before the pores became clogged. Intensive careful cleaning will help lengthen strainer use for additional 2-3 times, but new ones are preferred after pore clogging, and further cleaning will result in a hole in the strainer.*
13. Freshly prepared trichome batches (in water) and trichome-dislodged leaves (control) were immediately stored at 4 °C separately until all tissue sections were processed.
14. 1 ml of water containing trichomes was stained fresh by adding 0.5 ml of 0.4 % trypan blue. Contents were incubated for 3-4 h at room temperature (in the light or dark) without shaking in a petri-dish in preparation for microscopy.

15. An Axiovert 100 microscope (10x magnification) was used to confirm the proportion of un-ruptured trichome cells and the quality of the preparation.


*Notes:*

- a. *The trichome isolation protocol has no washing step, but to obtain the image of trichomes (Figure 1) we conducted a washing step using water for improving the appearance of the trichomes.*
- b. *Intact cells are unstained; disrupted cells are stained blue; a poor quality preparation would contain more disrupted trichomes and/or a greater proportion of contaminating leaf tissue fragments compared with Figure 1.*

16. The samples were then flash frozen in liquid N<sub>2</sub>, stored at -80 °C for 4-5 h, and freeze dried overnight in a Flexidry™ unit.

17. The method can be used for purifying non-glandular trichomes from other plant species (Figure 1).

18. Suggested leaf sample size for isolating *B. villosa* trichomes for various purposes (leaves with 300-500 µm trichome length, ~4,000 trichomes/cm<sup>2</sup> and 1 mg of dried trichomes isolation from 100 g of fresh leaf tissue [FL]): (a) for electron microscopic structural analysis (100/cm<sup>2</sup>); (b) for RNA isolation (300 g FL); (c) for biochemical extraction (500 g FL); for HPLC-UV analysis (500 g FL); for NMR structural analysis (800-1,000 g FL); for metal content analysis by inductively coupled plasma mass spectroscopy (500 g FL).



**Figure 1. High quality trichome preparations at 10x magnification after isolation from a *Brassica villosa* b Soybean (*Glycine max*) c Tomato (*Solanum lycopersicum*). Arrows indicate contaminating leaf tissue fragments or broken trichomes.**

## Representative data

### **Video 1. Detailed procedure for trichome isolation from *Brassica villosa***



## Recipes

1. 0.4 % trypan blue  
40 mg of trypan blue dissolved in 100 ml Millipore-filtered water

## Acknowledgments

This research was supported by grants from the Canola Council of Canada, Sask-Canola, the Saskatchewan Agriculture Development Fund, and funding from Agriculture and Agri-Food Canada.

## References

1. Aziz, N., Paiva, N. L., May, G. D. and Dixon, R. A. (2005). [Transcriptome analysis of alfalfa glandular trichomes](#). *Planta* 221(1): 28-38.
2. Konna, P. and Jackai, L. (2004). [The potential of pod-shaving in studies of the role of trichomes in \*Vigna\* resistance to the pod-bug \*Clavigralla tomentosicollis\* Stal \(Hemiptera: Coreidae\)](#). *International Journal of Tropical Insect Science* 24(04): 298-303.
3. Levin, D. A. (1973). [The role of trichomes in plant defense](#). *Q Rev Biol*: 3-15.
4. Mathur, J. (2006). [Trichome cell morphogenesis in \*Arabidopsis\*: a continuum of cellular decisions](#). This review is one of a selection of papers published in the Special Issue on Plant Cell Biology. *Botany* 84:604-612.

5. Nayidu, N. K., Tan, Y., Taheri, A., Li, X., Bjorndahl, T. C., Nowak, J., Wishart, D. S., Hegedus, D. and Gruber, M. Y. (2014). [\*Brassica villosa\*, a system for studying non-glandular trichomes and genes in the Brassicas.](#) *Plant Mol Biol* 85(4-5): 519-539.
6. Zhang, X. and Oppenheimer, D. G. (2004). [A simple and efficient method for isolating trichomes for downstream analyses.](#) *Plant Cell Physiol* 45(2): 221-224.