

## Small-scale Triton X-114 Extraction of Hydrophobic Proteins

Yuzuru Taguchi\* and Hermann M. Schätzl

Comparative Biology and Experimental Medicine, University of Calgary, Calgary, Canada

\*For correspondence: [ytaguchi@ucalgary.ca](mailto:ytaguchi@ucalgary.ca)

**[Abstract]** Here we introduce a protocol for Triton X-114 extraction which we used in our recently-published paper (Taguchi *et al.*, 2013). It is a versatile method to concentrate or partially purify hydrophobic proteins. The presented protocol is based on the protocol published by Bordier (Bordier, 1981) but more simplified and down-scaled for more small-scale and simpler use (Taguchi *et al.*, 2013).

Triton X-114 (TX114) is a non-ionic detergent which has a relatively low clouding point at 22 °C and separates into detergent (Det) and aqueous (Aq) phase at temperatures above the clouding point. During phase separation, hydrophobic solutes in the TX114 solution are sequestered to the Det phase, while hydrophilic solutes are sequestered to the Aq phase. Utilizing this phenomenon, TX114 extraction is a very versatile technique to efficiently concentrate hydrophobic proteins, especially glycosylphosphatidylinositol (GPI)-anchored proteins like the prion protein (PrP), because they have substantial amounts of highly hydrophobic moieties. Besides, phase separation using TX114 tolerates a variety of conditions, e.g. different pH or relatively low concentrations of guanidine hydrochloride. Since the hydrophobic proteins are sequestered to the Det phase as long as the phase separation occurs, and if the hydrophobicity of the protein of interest is not affected by pH or denaturant, this technique can be also utilized to change buffers or to remove denaturants. When using enzymes or proteases which maintain activities in detergent solutions, TX114 can also be used to separate hydrophobic from the water-soluble hydrophilic moieties upon enzymatic digestion of proteins, as done by us using *in vitro* digestion of PrP with phosphatidylinositol-specific phospholipase C (Taguchi *et al.*, 2013).

### **Materials and Reagents**

1. Neuro2a cells (N2a) (100%-confluent on a well of a 6-well plate)
2. Triton X-114 (Sigma-Aldrich, catalog number: X 114-100 ML)
3. 10x phosphate-buffered saline (Sigma-Aldrich, catalog number: P5493-1L)
4. Complete protease-inhibitor cocktail (Roche Diagnostics, catalog number: 04693116001)  
*Note: Optional, depending on the kind of the protein of interest and the purpose of experiments.*
5. 0.5 M EDTA (EMD Millipore, catalog number: 324503-1KG)

6. Deoxycholic acid (Sigma-Aldrich, catalog number: D2510-100G)
7. Triton X-100 (Sigma-Aldrich, catalog number: T8787-100ML)
8. Sodiumdodecyl sulfate (Sigma-Aldrich, catalog number: L6026-50G)
9. Glycerol (Sigma-Aldrich, catalog number: G9012-500ML)
10. Purified water
11. Phosphate-buffered saline without calcium and magnesium (Ca/Mg) (Life Technologies, catalog number: 10010-023)
12. Phosphate-buffered 0.5% Triton X-100 (TX100)/0.5% deoxycholate (DOC) lysis buffer (see Recipes)
13. 5x sample buffer (see Recipes)

### **Equipment**

1. Refrigerated centrifuge (max rcf 16,100 x g) (Eppendorf, model: 5415R)
2. Non-refrigerated centrifuge (max rcf 21,130 x g) (Eppendorf, model: 5424)
3. 50 ml conical tube
4. 6-well plate
5. 1.5 ml plastic tube

### **Procedure**

#### A. Pre-condensation

This is a step to remove hydrophilic components from the detergent so that the amounts of TX114 which unfavorably sequester to the Aq phase are reduced and, at the same time, the clouding point is lowered. Although dibutyl hydroxytoluene is used in the precondensation procedure in the original protocol, presumably as an antioxidant to prevent auto-oxidation of TX114, eliminating it from this step does not affect the efficacy of phase-separation. One point to note is that certain concentrations of electrolytes are indispensable for efficient phase separation of TX114. We usually use 1-10 mM Tris-HCl or phosphate buffer, depending on the following experiments. Described here is the specific protocol for our experiments. As this step takes hours, we recommend finishing this step at least by the previous day of harvesting the cells.

1. Place 47 ml of buffer in a graduated 50 ml conical tube, then add 3 ml of TX114. As TX114 is very viscous, repeat pipetting until most of the TX114 is homogenized in the buffer. Then, mix well by vortexing or rigorous shaking.
2. Clarify the solution on ice or 4 °C, then mix again. This usually takes a few minutes.
3. Incubate the tube at 37 °C to let the solution phase-separate by gravity. As viscous

greasy phase which is heavier than water, the Det phase is formed below the Aq phase. Since the concentration of TX114 in the Det phase after phase separation is ~12% (Bordier, 1981), the volume of the detergent phase is expected to be 24 ml. It usually takes a few hours for the top of the Det phase to descend to this level.

4. When the Det phase falls to the expected level, aspirate the Aq phase and discard it. Then, add fresh buffer up to 50 ml, mix well by vortexing and shaking, and clarify the solution on ice again.
5. Repeat the steps A3-4, 3-4 times. After the last cycle of phase-separation, discard the Aq phase and store in the dark. The precondensed TX114 can be stored at 4 °C in the dark for at least 6 months.

#### B. Preparation of TX114 lysis buffer and wash buffer

We usually prepare 2% TX114 lysis buffer. As the concentration of TX114 in the precondensed TX114 (detergent phase) is ~12%, we add one part of the precondensed TX114 to five parts of tris- or phosphate-buffered saline (PBS).

1. Take 37 ml of purified water in a 50 ml conical tube and add 5 ml of 10x PBS and 8 ml of the precondensed TX114. As viscosity of the solution is lower when the solution is turbid, this step is done at RT. Repeat pipetting until the TX114 sticking to inside the pipette is removed.
2. Clarify the lysis buffer on ice. If it has already phase-separated, mix well then clear it on ice again.
3. Take 2.5 ml of 2% TX114 lysis buffer and add to 47.5 ml of buffered saline, e.g. 5 ml of 10x PBS + 42.5 ml of purified water, to make "0.1% TX114 wash buffer".
4. Store the lysis buffer and the wash buffer at 4 °C.

#### C. Lysis of cells

As TX114 is a mild detergent, lysis by directly pouring the lysis buffer onto the cells on the plate is inefficient. Therefore, we usually first detach the cells with 3 mM EDTA in Ca/Mg-free PBS, collect the cells in a tube, then lyse them by addition of 2% TX114 lysis buffer.

1. Remove old culture medium from the N2a cells cultured on a well of a 6-well plate and rinse the monolayer of cells with Ca/Mg-free PBS.
2. Then, add 3 mM EDTA in Ca/Mg-free PBS (600 µl), and incubate for a few minutes until the cells can be easily detached by pipetting.

*Note: Be careful not to incubate too long. Although our N2a cells can be detached by this procedure, more adhesive cells might require cell scrapers for detachment.*

3. After completely detaching the cells by pipetting, collect the cell suspension in a 1.5 ml plastic tube. Rinse the bottom of the well with 400 µl of 3 mM EDTA in Ca/Mg-free PBS

and add it to the tube. Keep the cell suspension on ice all along.

4. Centrifuge the cell suspension at 1,000 x g, at 4 °C, for 5 min. Then, carefully aspirate and discard the supernatant.
5. Add 300 µl-1,000 µl of phosphate-buffered 2% TX114 lysis buffer and vortex rigorously for ~30 sec. The volume of the lysis buffer to be added depends on the purpose of experiments and the following procedures.
  - a. With larger volumes of lysis buffer, volumes of the Det phase after phase-separations are accordingly larger.
  - b. Incubate the cell suspension on ice for ~30 min with vortexing for ~5 sec from time to time.
6. Centrifuge the cell suspension at 16,100 x g for 2 min at 4 °C to precipitate the cell debris.
  - a. Make sure that the temperature of the rotor is low enough not to let the lysate phase-separate during the centrifugation.
  - b. Although the clouding point of pure TX114 solution is ~22 °C, it tends to be lower with higher concentrations of impurities.
7. Carefully take the supernatant to a new tube as the TX114 lysate.

#### D. Phase separation

1. Incubate the TX114 lysate at 37 °C for ~10 min until the lysate gets turbid and the Det phase partly precipitates.
2. Centrifuge at 22,500 x g for 10 min at room temperature. Make sure the rotor is warm enough for efficient phase-separation.
3. Gently take out the tube and carefully remove the Aq phase. If the hydrophilic proteins in the Aq phase are required for the experiments, put the Aq phase in a new tube.
4. Add 0.1% TX114 wash buffer and mix well by vortexing and/or shaking. We usually add the same volume as the eliminated Aq phase but different volumes should not cause problems. Then, clarify the lysate on ice.
5. Incubate the lysate at 37 °C for phase-separation. Repeat steps D1-4 2 more times.

#### E. After phase-separation

We usually precipitate the hydrophobic proteins extracted to the Det phase by methanol/chloroform precipitation (Taguchi *et al.*, 2013) and then add sample buffer and boil for preparation of the samples for SDS-PAGE and immunoblotting.

However, other usage of the partially purified hydrophobic proteins is possible, if the detergent can be efficiently removed.

## Recipes

1. Phosphate-buffered 0.5% Triton X-100 (TX100)/0.5% deoxycholate (DOC) lysis buffer
  - a. First prepare 5% TX100/DOC stock solution
 

Triton X-100	5 ml
Deoxycholic	5 g
Purified water	up to 100 ml
  - b. Then mix 5% TX100/5% DOC, 10x phosphate-buffered saline (PBS) and water in a 50 ml-conical tube
 

5% TX100/5% DOC	5 ml
10x PBS	5 ml
Purified water	up to 50 ml
2. 5x sample buffer
 

SDS	1.2 g
1 M Tris-HCl (pH 7.1)	2.5 ml
Glycerol	4 ml
0.5% BPB	300-500 µl
Water	up to 10 ml

## Acknowledgments

This protocol was adapted from Taguchi *et al.* (2013). This work was supported by grants for the National Institute of Health R01 NS076853-01A1 and the Alberta Prion Research Institute (AB, Canada).

## References

1. Bordier, C. (1981). [Phase separation of integral membrane proteins in Triton X-114 solution.](#) *J Biol Chem* 256(4): 1604-1607.
2. Taguchi, Y., Mistica, A. M., Kitamoto, T. and Schatzl, H. M. (2013). [Critical significance of the region between Helix 1 and 2 for efficient dominant-negative inhibition by conversion-incompetent prion protein.](#) *PLoS Pathog* 9(6): e1003466.