

Protein Degradation Assays in Arabidopsis Protoplasts

Filip Mituła^{1*}, Anna Kasprowicz-Maluśki², Michał Michalak², Małgorzata Marczak¹, Konrad Kuczyński¹
and Agnieszka Ludwików^{1*}

¹Department of Biotechnology, Institute of Molecular Biology and Biotechnology, Faculty of Biology, Adam Mickiewicz University in Poznań, Poznań, Poland; ²Department of Molecular and Cellular Biology, Institute of Molecular Biology and Biotechnology, Faculty of Biology, Adam Mickiewicz University in Poznań, Poznań, Poland

*For correspondence: mitula@amu.edu.pl and ludwika@amu.edu.pl

[Abstract] Plant transformation and exogenous protein expression is essential for molecular biology and biotechnology. Current approaches of stable plant transformation might be problematic and very time-consuming. Because of this, transient expression in protoplasts has become valuable alternative, being less cost and time-effective at the same time. Excellent for eukaryotic proteins, representing a natural cell habitat, protoplast isolation is widely used in protein interaction visualization techniques, like BiFC (Bimolecular fluorescence complementation) and FRET (Förster resonance energy transfer). In this protocol we present a another use of *Arabidopsis* protoplast in protein degradation assay, proving its high versatility as a tool in proteomics.

Materials and Reagents

1. 3-week old *Arabidopsis* plants
2. Mannitol (BDH Prolabo, catalog number: 25311)
3. CaCl₂ (POCH, catalog number: M00015143)
4. KCl (USB, catalog number: 20598)
5. 2-(N-morpholino)ethanesulfonic acid (MES) (LabEmpire, catalog number: MES503)
6. NaCl (POCH, catalog number: BA4121116)
7. Polyethylene glycol (PEG) 4000 (Sigma-Aldrich, catalog number: 81240)
8. Cellulase (SERVA Electrophoresis GmbH, catalog number: 16419)
9. Macerozyme (SERVA Electrophoresis GmbH, catalog number: 28302)
10. GenElute™ HP Plasmid Midiprep Kit (Sigma-Aldrich, catalog number: PLD35)
11. Benzyloxycarbonyl-L-leucyl-L-leucyl-L-leucinal, Z-Leu-Leu-Leu-al (MG132) (Sigma-Aldrich, catalog number: M7449)
12. Dimethyl sulfoxide (DMSO) (Sigma-Aldrich, catalog number: D8418)
13. HEPES buffer (Sigma-Aldrich, catalog number: H3375)
14. MgCl₂ (USB, catalog number: 18641)
15. Dithiothreitol (DTT) (Sigma-Aldrich, catalog number: D0632)
16. Phenylmethylsulfonyl fluoride (PMSF) (Sigma-Aldrich, catalog number: P7626)
17. Protease inhibitors (Roche Diagnostics, catalog number: 11873580001)

18. Enzyme solution (see Recipes)
19. W5 (see Recipes)
20. W1 (see Recipes)
21. MMg (see Recipes)
22. 40% PEG (see Recipes)
23. Protein isolation buffer (see Recipes)

Equipment

1. Light microscope (Nikon Corporation)
2. Hemocytometer (Sigma-Aldrich)
3. Tape (Scotch® Magic™)
4. Stainless steel forceps (Sigma-Aldrich)
5. Scissors
6. 10 ml pipette
7. 50 ml tubes
8. 90 mm Petri dishes
9. Amicon Ultra-15 centrifugal filter unit (Millipore)
10. NanoDrop spectrophotometer
11. Tabletop centrifuge
12. Horizontal shaker

Procedure

A. Protoplast isolation

1. Using tweezers, 5-6 mature leaves of *Arabidopsis*, petioles removed, were harvested.
2. Adaxial surface of leaves was put on a piece of tape and leaves were flattened (Figure 1A-B).
3. Second piece of tape was applied with limited pressure, trapping leaves between them (Figure 1C). Too much pressure will result in damaging leaves, too little pressure will lead to poor epidermis removal effectiveness.
4. Sandwich was placed with adaxial surface on top and pieces of tape were split by pulling away tape on top side, started at the tip of leaf (Figure 1D-E) (see Note 2).




Figure 1. Preparation of Arabidopsis leaf for protoplast isolation. A. 3-weeks old *Arabidopsis thaliana*; B-E. Step by step procedure of leaf epidermis removal with tape.

5. Tape was cut around leaves and put at room temperature enzyme solution in Petri dish, exposed mesophyll down. Tape should not be immersed in solution, but float on top of it. Mesophyll cells were digested for 60 min at 30 °C with gentle shaking (55 rpm) (see Note 3).
6. After digestion pieces of tape were removed and cells were left for another 5 min in same conditions (see Note 4).
7. Using 10 ml pipette protoplasts were transported to 50 ml tubes and put on ice.
8. Protoplast were centrifuged for 3 min (150 x g, 4 °C) and washed twice with W5 buffer. Be careful not to resuspend protoplast abruptly. After second wash step protoplast were resuspended in 1 ml of MMg solution. Protoplasts were calculated with help of hemocytometer and diluted to optimal concentration of 2×10^4 cells in 100 μ l with MMg solution (see Notes 5-7).
9. 5-10 μ g plasmid DNA coding a tag-protein fusion was aliquoted to 2 ml eppendorf tube (see Note 8).
10. 100 μ l of isolated protoplasts in MMg medium were transferred to the tube using a pipette tip with the tip of the tip cut off.
11. Using a pipette tip with the tip of the tip cut off 110 μ l of PEG solution was added. Solution was gently mixed and left for 15 min at RT in horizontal position (see Note 9).
12. After incubation, 450 μ l of W1 buffer was added to the tubes to dilute PEG solution, mixed and centrifuged for 3 min, 300 x g.
13. Supernatant was removed and harvested protoplasts were resuspended in 300 μ l of W1 solution.
14. Transfected protoplasts were incubated overnight at 22 °C in horizontal position in the dark for protein expression. It is important not to disturb protoplasts at this stage.
15. From this point protoplasts can be used in various protein analysis techniques (kinase assay, *in vivo* protein degradation assay, protein subcellular localization, BiFC and FRET analyses) (Ludwikow *et al.*, 2014). Here we show *in vivo* degradation assay for plant proteins.

B. *In vivo* degradation assay in protoplasts

16. After overnight incubation (as indicated in step A14) in the dark protoplasts were treated with 50 μ M MG132 (proteasome inhibitor) or mock treated with 0.1% DMSO for 6 h.
17. After brief centrifugation (300 x g, RT) supernatant was removed and cells were resuspended and disrupted in 100 μ l of protein isolation buffer.
18. Protein concentration was determined using a NanoDrop spectrophotometer.
19. Prepared samples were separated by SDS-PAGE and further analyzed by Western blotting.

Representative data




Figure 2. MG132 treatment increases ACC synthase6 protein accumulation in the *abi1td* protoplasts.

Protoplasts isolated from WT Col-0 and the *ABI1* knockout line (*abi1td*) were transformed with 5 µg of DNA plasmid coding for StrepTag-ACS6. Transformed protoplasts were treated with 50 µM MG132 or an equivalent volume of DMSO (mock control) for 6 h prior harvesting. A Western blot with anti-StrepTag antibodies confirms the presence of the StrepTag-ACS6 protein. The Western blot shown is representative of at least three independent experiments. Coomassie staining confirms equal protein loading (Ludwikow *et al.*, 2014).

Notes

1. This protocol is applicable to Brassica napus protoplast isolation and transformation.
2. To avoid mesophyll cells damage (visible as dark green spots) don't use too much pressure when applying the tape.
3. It is not recommended to digest leaves for more than 60 min. Protoplasts yield at this point will not increase, but they lose viability.
4. If one hour digestion did not freed all mesophyll cells from tape fragments, one can gentle dip tape a few times in enzyme solution, to increase protoplast yield.
5. Use swinging bucket rotor for centrifugation.
6. Keep low acceleration and deceleration values during centrifugation.
7. Resuspend the protoplasts by gently rocking the tube.
8. Use hemocytometer to achieve accurate and reproducible results.
9. A large volume of plasmid DNA decreases transformation efficiency, therefore keep the volume around 10 µl. Low transformation efficiency is usually a result of low quality plasmid DNA.

Recipes

1. Enzyme solution (10 ml)
1.2% cellulose

- 0.4% macerozyme
- 0.4 M mannitol
- 20 mM KCl
- 20 mM MES
- dH₂O
- Filter sterilize
- Incubate at 55 °C for 10 min
- Prepare fresh, do not store
- 2. W5 (50 ml)
 - 154 mM NaCl
 - 125 mM CaCl₂
 - 5 mM KCl
 - 2 mM MES
 - dH₂O
 - Filter sterilize, autoclave
 - Stored at 4 °C
- 3. W1 (10 ml)
 - 0.5 M mannitol
 - 20 mM KCl
 - 4 mM MES
 - dH₂O
 - Filter sterilize, autoclave
 - Stored at 4 °C, but no longer than 2 weeks
- 4. MMg (10 ml)
 - 0.4 M mannitol
 - 15 mM MgCl₂
 - 4 mM MES
 - dH₂O
 - Filter sterilize, autoclave
 - Stored at 4 °C, but no longer than 2 weeks
- 5. 40% PEG (10 ml)
 - 4 g PEG 4000
 - 200 mM mannitol
 - 100 mM CaCl₂
 - Filter sterilize, autoclave
 - Stored at 4 °C, but no longer than 3 weeks
- 6. Protein isolation buffer
 - 20 mM HEPES (pH 7.5)
 - 10 mM MgCl₂
 - 1 mM DTT

1 mM PMSF
Protease inhibitor
dH₂O
Stored at -20 °C

Acknowledgments

This work was supported by COST Action FA0605 project 682/N-COST/2010/0, the National Science Centre grants (5615/B/P01/2010/39, DEC-2012/05/B/NZ3/00352, DEC-2011/03/N/NZ3/01796) and POLAPGEN grant no. WND-POIG.01.03.01-00-101/08. This protocol was adapted from Wu *et al.* (2009).

References

1. Ludwików, A., Ciesła, A., Kasproicz-Maluśki, A., Mitula, F., Tajdel, M., Galganski, L., Ziolkowski, P. A., Kubiak, P., Malecka, A., Piechalak, A., Szabat, M., Gorska, A., Dabrowski, M., Ibragimow, I. and Sadowski, J. (2014). [Arabidopsis protein phosphatase 2C ABI1 interacts with type I ACC synthases and is involved in the regulation of ozone-induced ethylene biosynthesis](#). *Mol Plant* 7(6): 960-976.
2. Wu, F. H., Shen, S. C., Lee, L. Y., Lee, S. H., Chan, M. T. and Lin, C. S. (2009). [Tape-Arabidopsis Sandwich - a simpler Arabidopsis protoplast isolation method](#). *Plant Methods* 5: 16.