


## LC/MS-based Detection of Hydroxyproline O-galactosyltransferase Activity

Mari Ogawa-Ohnishi and Yoshikatsu Matsubayashi\*

Division of Biological Science, Graduate School of Science, Nagoya University, Nagoya, Japan

\*For correspondence: [matsu@bio.nagoya-u.ac.jp](mailto:matsu@bio.nagoya-u.ac.jp)

**[Abstract]** Arabinogalactan proteins (AGPs) are plant-specific extracellular glycoproteins regulating a variety of processes during growth and development. AGP biosynthesis involves O-galactosylation of hydroxyproline (Hyp) residues followed by a stepwise elongation of the complex sugar chains. The initial Hyp O-galactosylation is mediated by Hyp O-galactosyltransferase (HPGT) that catalyzes the transfer of a D-galactopyranosyl residue to the hydroxyl group of Hyp residues of peptides from the sugar donor UDP- $\alpha$ -D-galactose (Figure 1). Here we describe a LC/MS-based method for the detection of HPGT activity *in vitro*.



**Figure 1. Reaction scheme for Hyp galactosylation by HPGT.** HPGT catalyzes the addition of a D-galactopyranose from an UDP- $\alpha$ -D-Gal to the hydroxylgroup of Hyp residues.

### Materials and Reagents

1. 1-week-old *Arabidopsis* T-87 cells (50 g fresh weight)
2. Bio-Rad Protein Assay (Bio-Rad Laboratories, catalog number: 5000006JA)
3. 2 mM synthesized substrate peptide [e.g., (OAOSOT)<sub>3</sub>S] [using standard Fmoc solid-phase synthesis chemistry on a 431A peptide synthesizer (Life Technologies)]
4. 2 mM Uridine 5'-diphosphogalactose disodium salt (Sigma-Aldrich, catalog number: U4500)
5. 1 M MOPS-KOH (pH 7.0)
6. 10 mM MnCl<sub>2</sub>
7. 10% TX-100
8. 1% Formic acid
9. Acetonitrile (HPLC grade) containing 0.1% formic acid
10. Water (HPLC grade) containing 0.1% formic acid

11. Tris-HCl (pH 7.0)
12. MgCl<sub>2</sub>
13. Dithiothreitol
14. Leupeptin
15. Phenylmethanesulfonyl fluoride
16. Sucrose
17. Extraction buffer (see Recipes)
18. Suspension buffer (see Recipes)

### **Equipment**

1. Waring blender
2. Miracloth (Merck Millipore Corporation, catalog number: 475855)
3. Ultracentrifuge
4. 30 °C incubator
5. Micro centrifuge
6. Semi-micro HPLC system (JASCO International Co., model: Micro21LC)
7. LCQ Deca XP-plus ESI ion-trap mass spectrometer (Thermo Fisher Scientific)
8. TSK-gel Amide-80 (3 μm) column (2 x 150 mm) (Tosoh Bioscience LLC, catalog number: 21865)

### **Procedure**

#### A. Preparation of *Arabidopsis* microsomal membranes

1. *Arabidopsis* T-87 cells are maintained on a 1-week culture interval under continuous darkness at 22 °C with shaking at 120 rpm.
2. Suspend 1-week-old *Arabidopsis* T-87 cell (50 g fresh weight) in 20 ml Extraction buffer.
3. Cool off to 4 °C on ice.
4. Homogenize at 20,000 rpm for 5 min at 4 °C in a Waring blender.
5. Cool off to 4 °C on ice.
6. Filter the slurry through two layers of Miracloth.
7. Centrifuge the filtrate at 3,000 x g for 15 min at 4 °C.
8. Centrifuge the supernatant at 100,000 x g for 30 min at 4 °C.
9. Suspend the pellet (microsomal membranes: approximately 150 μg/μl) in 500 μl Suspension buffer by gentle pipetting.
10. Determine the protein concentration by conventional Bradford assay according to the manufacturer's protocol (Bio-Rad Protein Assay).

B. Hyp O-galactosyltransferase activity assay

1. Set up 20  $\mu$ l HPGT assay reactions in 0.5 ml microcentrifuge tube as follows.


HPGT assay components	Amount per reaction
1 M MOPS-KOH (pH 7.0)	2 $\mu$ l
2 mM UDP- $\alpha$ -D-galactose	5 $\mu$ l
10 mM MnCl <sub>2</sub>	2 $\mu$ l
10% TX-100	2 $\mu$ l
2 mM substrate peptide	1 $\mu$ l
<i>Arabidopsis</i> T-87 microsomal membranes	30 $\mu$ g protein equivalent
Water	Total 20 $\mu$ l

2. Incubate at 30 °C, 1 h.
3. Add 2  $\mu$ l 1% formic acid to stop reaction.
4. Add 80  $\mu$ l acetonitrile.
5. Centrifuge at 20,000 x g for 5 min.

C. LC/MS analysis

10  $\mu$ l aliquots of the assay solution were analyzed by LC-MS using a micro HPLC (high pressure liquid chromatography) system connected to an LCQ Deca XP-plus ESI ion-trap mass spectrometer. Chromatographic separation is performed by normal-phase HPLC on a TSK-gel Amide-80 (3  $\mu$ m) column (2 x 150 mm).

1. The mobile phase is composed of HPLC grade water containing 0.1% formic acid (eluent A) and HPLC grade acetonitrile containing 0.1% formic acid (eluent B). The column temperature is maintained at 25 °C.
2. The HPLC flow rate is 100  $\mu$ l/min, and the elution gradient was 60 to 30% B over 15 min.
3. The HPLC eluate was introduced into an electrospray ionization (ESI) ion-trap mass spectrometer in the positive ionization mode.
4. MS source parameters are as follows [e.g., (OAOSOT)<sub>3</sub>S peptide]:
  - a. Capillary temperature: 200 °C
  - b. Capillary voltage: 42 V
  - c. Source voltage: 5 kV
  - d. Source current: 8.5  $\mu$ A
  - e. Sheath gas flow: 50
  - f. Aux gas flow: 0
  - g. Sweep gas flow: 0
  - h. The mass range: *m/z* 500-2,000
5. The mass spectra are obtained by selected ion monitoring. [e.g., (OAOSOT)<sub>3</sub>S: *m/z* 951.1, Galactosylated product: *m/z* 1032.2] (Figure 2).



**Figure 2. Selected ion chromatogram of substrate peptide and the galactosylated product.** Substrate peptide was incubated with solubilized membrane fractions in the presence of UDP- $\alpha$ -D-galactose, then analyzed by LC-MS with selected ion monitoring of the substrate ( $m/z$  951.1) and the galactosylated product ( $m/z$  1032.2).

### Recipes

1. Extraction buffer (prepare freshly and keep on ice)
  - 25 mM Tris-HCl (pH 7.0)
  - 10 mM MgCl<sub>2</sub>
  - 2 mM dithiothreitol
  - 2  $\mu$ M leupeptin
  - 2 mM phenylmethanesulfonyl fluoride
  - 250 mM sucrose
2. Suspension buffer
  - 10 mM Tris-HCl (pH 7.0)
  - 250 mM sucrose

### **Acknowledgments**

This is the detailed protocol for the detection of HPGT activity described by Ogawa-Ohnishi and Matsubayashi (2015). This research was supported by a Grant-in-Aid for Scientific Research (S) from the Ministry of Education, Culture, Sports, Science, and Technology (No. 25221105).

### **References**

1. Ogawa-Ohnishi, M. and Matsubayashi, Y. (2015). [Identification of three potent hydroxyproline O-galactosyltransferases in Arabidopsis](#). *Plant J* 81(5): 736-746.
2. Ogawa-Ohnishi, M., Matsushita, W. and Matsubayashi, Y. (2013). [Identification of three hydroxyproline O-arabinosyltransferases in Arabidopsis thaliana](#). *Nat Chem Biol* 9(11): 726-730.