

## Measurement of NADPH Oxidase Activity in Plants

Amita Kaundal<sup>1</sup>, Clemencia M. Rojas<sup>1</sup> and Kirankumar S. Mysore<sup>2\*</sup>

<sup>1</sup>Plant Biology Department, The Samuel Roberts Noble Foundation, Ardmore, USA;

\*For correspondence: [kmysore@noble.org](mailto:kmysore@noble.org)

**[Abstract]** NADPH oxidase is a membrane-bound enzyme that generates ( $O_2^{\cdot-}$ ) by transferring electrons from NADPH to molecular oxygen  $O_2$ .  $O_2^{\cdot-}$  is spontaneously dismasted to the more stable form  $H_2O_2$ . Both  $O_2^{\cdot-}$  and  $H_2O_2$  are forms of reactive oxygen species (ROS), which are involved in regulation of many cellular activities such as transcription, intracellular signaling, and host defense. The NADPH oxidase - dependent generation of  $O_2^{\cdot-}$  in total membrane fraction of plant tissue has been determined by the reduction of the tetrazolium salt XTT by  $O_2^{\cdot-}$ . In the presence of  $O_2^{\cdot-}$ , XTT generates a soluble yellow formazan that can be quantified spectrophotometrically.

### Materials and Reagents

1. Sucrose
2. HEPES
3. EDTA
4. DTT
5. L-cysteine
6.  $MgCl_2$
7. PVP
8. Complete, Mini, EDTA-free Protease Inhibitor Cocktail Tablets (F. Hoffmann-La Roche, catalog number: 04693159001)
9. BSA
10. Bio-Rad Protein Assay (Bio-Rad Laboratories, catalog number: 500-0006)
11. Tris-HCl
12. Sodium 3,3'-([[(phenylamino)carbonyl]-3,4-tetrazolium)-bis (4-methoxy-6-nitro) benzene-sulfonic acid hydrate (XTT) (Sigma-Aldrich, catalog number: X4626)
13. NADPH (Sigma-Aldrich, catalog number: N1630)
14. Protein extraction working solution (see Recipes)

### Equipment

1. Microtiter plate reader (Infinite M200 Pro, Tecan)

2. Microcentrifuge (AqquSpin Micro R) (Thermo Fisher Scientific)
3. Ultracentrifuge (Optima TLX, Beckman)
4. Microtiter plate (BD Biosciences, catalog number: 353075)

**Procedure**

- A. Protein extraction and separation of membrane fraction from plant tissues
  1. Harvest tissue in liquid nitrogen. If not used immediately, keep at -80 °C until processing.
  2. Grind tissue in liquid nitrogen and weigh out 0.5 g of the ground tissues in empty Falcon tube that has been pre-chilled in liquid nitrogen and used to tare the scale.
  3. Add 6 ml of ice-cold protein extraction buffer to ground tissues on ice.
  4. Vortex at room temperature to mix thoroughly.
  5. Filter homogenized tissue through four layers of cheese cloth and transfer filtrate (flow-through) to 2-ml microcentrifuge tubes, on ice.
  6. Centrifuge at 10,000 x g for 45 min at 4 °C and transfer supernatant to ultra- centrifuge tube.
  7. Separate total membrane fractions by ultra-centrifugation at 203,000 x g for 60 min at 4 °C.
  8. Discard supernatant and resuspend pellet in 1 ml ice-cold 10 mM Tris-HCl (pH 7.4).
  
- B. Protein estimation using Bradford microassay (160 µl)
  9. Prepare BSA standards ranging from 5 µg-25 µg/ml as follows:

Standard concentration	Volume of BSA (100µg/ml)	Volume of water
5 µg/ml	8 µl	152 µl
10 µg/ml	16 µl	144 µl
20 µg/ml	32 µl	128 µl
25 µg/ml	40 µl	120 µl

These standards will be used to generate a standard curve.

10. Use 96 well microtiter plate to prepare reaction mix.
11. Prepare blank by adding 160 µl of water to one well in triplicates.
12. Prepare test samples by adding 2 µl of supernatant (from section 1) to 158 µl of water
13. Add 40 µl of Bradford Assay reagent to BSA standards, blank and test samples.
14. Mix and incubate at room temperature for 5 min and read absorbance at 595 nm ( $A_{595}$ ) on plate reader spectrophotometer.

*Note: If spectrophotometer does not include a software to generate standard curve to automatically estimate protein content, generate a BSA standard curve by plotting known*

protein concentration (X-axis) vs. Absorbance (in Y-axis). Protein concentration for a given unknown sample is estimated by plotting the  $A_{595}$  absorbance of the unknown (in the y-axis) and determining the intersection point with the BSA standard curve and then find the concentration associated with that particular point (in the x-axis). If using excel, after plotting concentration vs  $A_{595}$ , obtain the trendline and use the equation for the line and the  $A_{595}$  of the unknown to resolve the unknown concentration.

C. NADPH oxidase activity assay

15. Prepare fresh solution of 1 mM XTT and 1 mM NADPH.
16. Prepare two different assay solutions A and B:

Stock solution	Assay solution A	Assay solution B
1M Tris-HCl (pH 7.5)	12.5 $\mu$ l	12.5 $\mu$ l
1mM XTT	125 $\mu$ l	125 $\mu$ l
1mm NADPH	25 $\mu$ l	-----
Water	77.5 $\mu$ l	102.5 $\mu$ l

*Note: Because membrane fraction can spontaneously reduce XTT, even in the absence of substrate (NADPH), it is necessary to prepare two blanks, one without NADPH, to correct for this background levels of activity.*

17. Prepare blanks:
  - Blank 1: Add 10  $\mu$ l of water to 240  $\mu$ l of Assay solution A.
  - Blank 2: Add 10  $\mu$ l of membrane fraction to 240  $\mu$ l Assay solution B.
18. Prepare samples by adding 10  $\mu$ l of membrane fraction (from section 1) to 240  $\mu$ l of Assay solution A. Read the absorbance at 492 nm ( $A_{492}$ ) at 0 min and then 20 min intervals for one hour or until saturation point reached.
19. To get final  $A_{492}$  Blank reading, subtracts  $A_{492}$  Blank1 and  $A_{492}$  Blank 2.
20. Calculate rate of  $O^{2-}$  generation by using an extinction coefficient  $2.16 \times 10^4 \text{ cm}$  (Jiang and Zhang 2002).

$$(\Delta A_{492\text{nm}}/\text{min}_{\text{test}} - \Delta A_{492\text{nm}}/\text{min}_{\text{blank}}) / (2.16 \times 10^4 \text{ M}^{-1} \text{ C}^{-1}) (0.04)$$

$$\Delta A_{492\text{nm}}/\text{min}_{\text{Test}} = A_{492\text{nm}}(\text{sample X}) \text{ at saturation point} - A_{492\text{nm}}(\text{sample X}) \text{ at 0 min}$$

$$\Delta A_{492\text{nm}}/\text{min}_{\text{blank}} = A_{492\text{nm}}(\text{blank}) \text{ at saturation point} - A_{492\text{nm}}(\text{blank}) \text{ at 0 min}$$

0.04 = dilution factor (10  $\mu$ l/250  $\mu$ l)

To calculate specific activity, divide the value obtained in equation by the amount of protein present in the sample (converted to mg/ml).

## Recipes

### 1. Protein extraction working solution

0.25 M sucrose

50 mM HEPES

3 mM EDTA

1 mM DTT

3.6 mM L-cysteine

0.1 mM MgCl<sub>2</sub>

0.6% PVP

10 Tablets of Complete, Mini, EDTA-free Protease Inhibitor Cocktail Tablets.

Prepare the following stock solutions:

1 M sucrose

1 M HEPES (pH 7.2)

0.25 M EDTA

1 M DTT

100 mM MgCl<sub>2</sub>

In 80 ml of water add the following reagents:

Stock solutions	Volume
1M Sucrose	25ml
1M HEPES (pH 7.2)	5ml
0.25M EDTA	1.2 ml
1M DTT	0.1 ml
L-cysteine	0.0632 g
100mM MgCl <sub>2</sub>	0.1 ml
PVP	0.6 g

Add 10 Tablets of Complete, Mini, EDTA-free Protease Inhibitor Cocktail Tablets.

Mix well and adjust volume to 100 ml

## Acknowledgments

This protocol has been adapted and modified to use in Arabidopsis from Jiang and Zhang (2002). This work was supported by the Samuel Roberts Noble Foundation.

---

**References**

1. Able, A. J., Guest, D. I. and Sutherland, M. W. (1998). [Use of a new tetrazolium-based assay to study the production of superoxide radicals by tobacco cell cultures challenged with avirulent zoospores of phytophthora parasitica var \*Nicotianae\*](#). *Plant Physiol* 117(2): 491-499.
2. Jiang, M. and Zhang, J. (2002). [Involvement of plasma-membrane NADPH oxidase in abscisic acid- and water stress-induced antioxidant defense in leaves of maize seedlings](#). *Planta* 215(6): 1022-1030.
3. Rojas, C. M., Senthil-Kumar, M., Wang, K., Ryu, C. M., Kaundal, A. and Mysore, K. S. (2012). [Glycolate oxidase modulates reactive oxygen species-mediated signal transduction during nonhost resistance in \*Nicotiana benthamiana\* and \*Arabidopsis\*](#). *Plant Cell* 24(1): 336-352.
4. Sagi, M. and Fluhr, R. (2001). [Superoxide production by plant homologues of the gp91\(phox\) NADPH oxidase. Modulation of activity by calcium and by \*Tobacco mosaic virus\* infection](#). *Plant Physiol* 126(3): 1281-1290.