

## Detection of D-glutamate Production from the Dual Function Enzyme, 4-amino-4-deoxychorismate Lyase/D-amino Acid Transaminase, in *Mycobacterium smegmatis*

Helen K. Opel-Reading, Roman Mortuza and Kurt L. Krause\*

Department of Biochemistry, University of Otago, Dunedin, New Zealand

\*For correspondence: [kurt.krause@otago.ac.nz](mailto:kurt.krause@otago.ac.nz)

**[Abstract]** D-amino acid transaminase (D-AAT) is able to synthesize both D-glutamate and D-alanine, according to the following reaction: D-alanine +  $\alpha$ -ketoglutarate = D-glutamate + pyruvate. These two D-amino acids are essential components of the peptidoglycan layer of bacteria. In our recently published work, MSMEG\_5795 from *Mycobacterium smegmatis* was identified as having D-amino acid transaminase (D-AAT) activity, although it has primarily been annotated as 4-amino-4-deoxychorismate lyase (ADCL). To unequivocally demonstrate D-AAT activity from MSMEG\_5795 protein two coupled enzyme assays were performed in series. First, D-alanine and  $\alpha$ -ketoglutarate were converted to D-glutamate and pyruvate by MSMEG\_5795 using the D-AAT assay. Next, the products of this reaction, following removal of all protein, were used as input into an assay for glutamate racemase in which D-glutamate is converted to L-glutamate by glutamate racemase (Gallo and Knowles, 1993; Poen *et al.*, 2016). As the only source of D-glutamate in this assay would be from the reaction of D-alanine with MSMEG\_5795, positive results from this assay would confirm the D-AAT activity of MSMEG\_5795 and of any enzyme tested in this manner.

**Keywords:** D-amino acid transaminase, Enzyme assay, Glutamate racemase, Transaminase, Amino-deoxychorismate lyase, Pyridoxal phosphate

**[Background]** The protocol described here in detail was developed during a study of suppressor mutants in *M. smegmatis* strain in which glutamate racemase (*murl*) had been deleted (Mortuza *et al.*, 2018). During this work, it became apparent that a mutation in the promoter of 4-amino-4-deoxychorismate lyase (ADCL) (MSMEG\_5795) was unexpectedly able to complement the *murl* deletion. This mutation resulted in a more than 10-fold increased expression of MSMEG\_5795. Because ADCL is homologous to D-amino acid transaminase (D-AAT), we decided to test MSMEG\_5795 for D-AAT activity. The initial test which involved incubation of the enzyme with D-alanine and  $\alpha$ -ketoglutarate was positive for activity. To be doubly sure that this protein, which was predominantly annotated as ADCL, was actually producing glutamate, we used the output of the D-AAT assay as input to the previously reported coupled assay for Murl (Tanizawa *et al.*, 1987; Gallo and Knowles, 1993). This assay is detailed below.

## **Materials and Reagents**

1. Pipette tips (Labcon, 1,250  $\mu$ l, catalog number: 1045-260-300; 200  $\mu$ l, catalog number: 1030-260-300; 10  $\mu$ l, catalog number: 1036-260-000)
  2. Microcentrifuge tubes 1.7 ml (MultiMax, catalog number: 2942)
  3. 50 ml conical centrifuge tubes (Cellstar, catalog number: 227261)
  4. Cuvette Type 9 quartz 10 mm path (Starna, catalog number: 9Q10)
  5. Vivaspin™ 2 protein concentrator spin column, MWCO 10,000 Da (GE Healthcare Life Sciences, catalog number: 28932247)
  6. Syringe filter, 0.45  $\mu$ m (Ahlstrom, catalog number: 760506)
  7. Syringe with Luer-Lok tip, 20 ml (Becton Dickinson, catalog number: 3028360)
  8. 10-15 mg/ml of purified *Mycobacterium smegmatis* MSMEG\_5795 in 20 mM Tris-HCl pH 8.0 and 150 mM NaCl (Mortuza *et al.*, 2018), and 20-40 mg/ml of purified *Bacillus anthracis* glutamate racemase isoform 2 (Murl2<sub>Ba</sub>) protein in 50 mM Tris pH 8.0, 250 mM NaCl and 0.1 mM DTT (May *et al.*, 2007)
  9. Tris base (NeoFroxx, catalog number: 1125KG001)
  10. HCl, 37%, reagent grade, ACS (Scharlau, catalog number: AC07412500)
  11.  $\alpha$ -ketoglutaric acid sodium salt (Sigma-Aldrich, catalog number: K1875)
  12. Pyridoxal 5'-phosphate hydrate (PLP) (Sigma-Aldrich, catalog number: P9255)
  13. L-lactate dehydrogenase from hog muscle (Roche, catalog number: 10107085001)
  14.  $\beta$ -Nicotinamide adenine dinucleotide, reduced dipotassium salt (NADH) (Sigma-Aldrich, catalog number: N4505)
  15. D-alanine (Sigma-Aldrich, catalog number: A7377)
  16. NaOH (ACS reagent grade, catalog number: SO042500)
  17. N-Cyclohexyl-2-aminoethanesulfonic acid (CHES) (ACROS, catalog number: 208181000)
  18. Iodonitrotetrazolium chloride (INT) (Sigma-Aldrich, catalog number: I10406)
  19. Diaphorase (Sigma-Aldrich, catalog number: D5540)
  20. Nicotinamide adenine dinucleotide (NAD<sup>+</sup>) (Sigma-Aldrich, catalog number: N1511)
  21. Adenosine 5'-diphosphate (ADP) (Sigma-Aldrich, catalog number: A2754)
  22. L-glutamate dehydrogenase (LGDH) (Sigma-Aldrich, catalog number: G2501)
  23. Pierce™ Coomassie Plus (Bradford) Assay Kit (Thermo Fisher, catalog number: 23236)
  24. 10 M NaOH (made by gradually dissolving NaOH pellets in ASTM Type I (18 M $\Omega$ ) water, stored at room temperature, no need to sterilize)
- Note: All solutions below are prepared in ASTM Type I (18 M $\Omega$ ) water and filtered through 0.45  $\mu$ m filters prior to use. Solutions are stored long-term at 4 °C, unless otherwise noted.*
25. 1 M Tris base (adjust the solution to pH 8.1 with HCl)
  26. 0.5 M  $\alpha$ -ketoglutaric acid
  27. 2 mM PLP
  28. 10 mM NADH (prepare this solution fresh for each assay)

29. 0.5 M D-alanine
30. 1 M CHES (adjust the solution to pH 9.2 with NaOH)
31. D-amino acid transaminase reaction partial mixture (see Recipes)
32. Glutamate reaction partial mixture (see Recipes)

## **Equipment**

1. Magnetic stir bars 15 mm x 6 mm (BRAND, catalog number: 137114)
2. Pipetman Neo<sup>®</sup> Pipets (Gilson, model: P1000N, catalog number: F144566; P200N, catalog number: F144565; P20N, catalog number: F144563)
3. 25 ml glass beaker (Boeco, catalog number: BOE 5010614)
4. Microcentrifuge (Eppendorf, model: 5415R)
5. Magnetic stirrer (Barnstead Thermolyne, model: SP131010-33)
6. Ultrospec<sup>™</sup> 3100 pro UV/Visible Spectrophotometer with SWIFT II software (GE Healthcare Life Sciences, model: Ultrospec<sup>™</sup> 3100 pro)
7. MultiTemp III (GE Healthcare) thermostatic circulator to control the temperature of the UV-Vis spectrophotometer cuvettes
8. -20 °C freezer

## **Software**

1. Biochrom<sup>™</sup> SWIFT II Software, version 2.05

## **Procedure**

### A. Assay Preparations

1. Purify 10-15 mg/ml MSMEG\_5795 and 20-40 mg/ml Murl2<sub>Ba</sub> protein using immobilized metal affinity chromatography and size exclusion chromatography.  
*Note: Determine the mycobacterial protein concentration using the Pierce<sup>™</sup> Coomassie Plus (Bradford) Assay according to the manufacturer's recommendation. Centrifuge both proteins at 16,000 x g for 20 min at 4 °C to remove any aggregates prior to being assayed.*
2. Prepare all stock solutions for the assays (see Recipes).

### B. D-amino acid transaminase reaction

1. Prepare desired volume of the D-AAT assay reaction partial mixture containing 100 mM Tris-HCl pH 8.1, 10 mM α-ketoglutarate, 0.15 mM PLP, 5 U L-lactate dehydrogenase, 0.2 mM NADH and 77 nM MSMEG\_5795 protein. For example, 20 ml is a convenient amount to make here. As noted above the NADH is prepared fresh each day before conducting the assay.  
*Note: Keep assay mix on ice during assay and equilibrate to room temperature before use.*

2. Transfer 950  $\mu$ l of the reaction mixture to a 1 ml quartz cuvette and pre-incubate at 30 °C for 5 min.
3. Measure the absorbance at 340 nm for 2 min to set a baseline on the Ultrospec™ 3100 pro.
4. To start the reaction, add 50  $\mu$ l 0.5 M D-alanine substrate to the assay mix to a final concentration of 25 mM to start the reaction and incubate each cuvette for 20, 40 and 60 min at 30 °C. The absorbance should be decreasing with time.
5. Add 50  $\mu$ l water instead of substrate to the negative control sample and incubate for 60 min at 30 °C.
6. Following incubation, transfer the cuvettes immediately to 4 °C to halt the reaction.
7. Filter the reaction mix using a Vivaspin 2 protein concentrator spin column with MWCO of 10,000 Da to remove MSMEG\_5795 and L-lactate dehydrogenase. Use this protein-free output as a substrate for the subsequent glutamate racemase (Murl) activity assay.

*Note: Samples can be stored at -20 °C before second coupled enzyme assay is performed or for further analysis.*

#### C. Glutamate racemase reaction

1. Prepare desired volume of Murl assay reaction partial mixture containing 50 mM CHES pH 9.2, 5 mM NAD<sup>+</sup>, 37.5 U ml<sup>-1</sup> L-glutamate dehydrogenase, 2.5 mM ADP, 0.65 mM INT, 2 U ml<sup>-1</sup> diaphorase. For example, 10 ml is a convenient amount to make here.

*Note: Mix assay components vigorously in a glass beaker, but avoid foaming by using a small stir bar and magnetic stirrer. Filter mixture with a 0.45  $\mu$ m syringe filter and a 20 ml syringe with a Luer-Lok tip. Keep solution on ice during assay. Equilibrate to room temperature before use. This solution will take on a light salmon hue.*


2. For each reaction sample, transfer 875  $\mu$ l of the reaction mixture to a 1 ml quartz cuvette.
3. Add 100  $\mu$ l of the product coming from the D-AAT assay to each cuvette.
4. Start the reaction by adding 25  $\mu$ l Murl2<sub>Ba</sub> to a final enzyme concentration of 10  $\mu$ M.
5. To the blank sample, add the buffer used for the Murl2<sub>Ba</sub> stock solution in place of Murl2<sub>Ba</sub> and add water instead of substrate.
6. Add water instead of substrate to the negative control sample.
7. Monitor the absorbance at 500 nm for 30 min at 30 °C.

*Note: Subtract the absorbance values obtained from the blank sample from each measurement to obtain the final absorbance values.*

#### Data analysis

The product of the D-AAT assay is taken as input for the glutamate racemase assay to confirm the presence of glutamate by monitoring the absorbance increase at 500 nm. The increase in absorbance values for the 20, 40 and 60 min samples corresponds to the expected increase in glutamate production over time (Figure 1). The detection of glutamate as reported here is qualitative.

This result was also supported with data obtained from high-resolution mass spectrometry of the assay output which showed an increase in glutamate concentration of 0.3, 0.7 and 1.0 mM in samples incubated for 20, 40 and 60 minutes respectively. As a result, the ability to convert D-alanine to D-glutamate by the MSEM<sub>G</sub>\_5795 enzyme can be assessed using the glutamate racemase assay.



**Figure 1. Time-dependent glutamate production by MSMEG\_5795 verified with glutamate racemase enzyme assay** (see Step C7). Protein-free product from the D-AAT assay was used as substrate for the glutamate racemase assay in the D to L direction. The increase in absorbance was observed at 500 nm for 30 min. Each sample was taken in duplicate and contained 100  $\mu$ l substrate and 10  $\mu$ M glutamate racemase. Blank sample was subtracted from each measurement and absorbance values were averaged before plotting against time. Absorbance values obtained from the 20, 40 and 60 min samples are shown in green, blue and red respectively. The control sample containing water instead of substrate is shown in black. Figure 1 is originally published in Mortuza *et al.*, 2018. (This figure is reprinted with permission from Molecular Microbiology.)

## Notes

1. The concentrations of the components listed in the assay mixtures represent the desired concentrations to have in place at the start of the assay, but be sure to take into account any dilution that may result from the addition of enzyme or other substance added subsequently.
2. If there is any concern that L-glutamate might be produced by the first enzyme being tested in this assay an additional control can be added in which the output of the first enzyme is assayed using the enzyme L-glutamate dehydrogenase directly without incubation with Murl.
3. Other glutamate racemases that are sufficiently active at 30 °C and pH 9.2 would likely work here as well, but we have only trialed this assay with the Murl<sub>2Ba</sub> enzyme.

## **Recipes**

1. D-amino acid transaminase reaction partial mixture
  - 100 mM Tris-HCl, pH 8.1
  - 10 mM  $\alpha$ -ketoglutarate
  - 0.15 mM PLP
  - 5 U L-lactate dehydrogenase
  - 0.2 mM NADH
  - 77 nM MSMEG\_5795 protein
  - Prepare in ASTM Type I (18 M $\Omega$ ) water
  - Filter through a 0.45  $\mu$ m filter prior to use
  - Long-term storage not recommended. Prepare fresh each day
  - Substrate is added later
2. Glutamate reaction partial mixture
  - 50 mM CHES, pH 9.2
  - 5 mM NAD<sup>+</sup>
  - 37.5 U ml<sup>-1</sup> L-glutamate dehydrogenase
  - 2.5 mM ADP
  - 0.65 mM iodinitrotetrazolium chloride
  - 2 U ml<sup>-1</sup> diaphorase
  - Prepare in ASTM Type I (18 M $\Omega$ ) water
  - Filter through a 0.45  $\mu$ m filter prior to use
  - Long-term storage not recommended. Prepare fresh each day
  - Substrate and Murl are added later

## **Acknowledgments**

This protocol was first described in brief in Mortuza *et al.* (2018). This research was supported by Lottery Health Research New Zealand-LHR2017-48905, the University of Otago, the Maurice Wilkins Centre-University of Auckland and the Thrash Foundation, Houston, Texas.

## **Competing interests**

The authors have no conflicts to declare.

## **References**

1. Gallo, K. A. and Knowles, J. R. (1993). [Purification, cloning, and cofactor independence of glutamate racemase from \*Lactobacillus\*](#). *Biochemistry* 32(15): 3981-3990.

2. May, M., Mehboob, S., Mulhearn, D. C., Wang, Z., Yu, H., Thatcher, G. R., Santarsiero, B. D., Johnson, M. E. and Mesecar, A. D. (2007). [Structural and functional analysis of two glutamate racemase isozymes from \*Bacillus anthracis\* and implications for inhibitor design](#). *J Mol Biol* 371(5): 1219-1237.
3. Mortuza, R., Aung, H. L., Taiaroa, G., Opel-Reading, H. K., Kleffmann, T., Cook, G. M. and Krause, K. L. (2018). [Overexpression of a newly identified D-amino acid transaminase in \*Mycobacterium smegmatis\* complements glutamate racemase deletion](#). *Mol Microbiol* 107(2): 198-213.
4. Poen, S., Nakatani, Y., Opel-Reading, H. K., Lasse, M., Dobson, R. C. and Krause, K. L. (2016). [Exploring the structure of glutamate racemase from \*Mycobacterium tuberculosis\* as a template for anti-mycobacterial drug discovery](#). *Biochem J* 473(9): 1267-1280.
5. Tanizawa, K., Masu, Y., Asano, S., Tanaka, H. and Soda, K. (1987). [D-Amino acid aminotransferase from a thermophile, YM-1: enzymological properties, cloning of the amino acid sequence](#). In: *The 7th International Congress on Chemical and Biological Aspects of Vitamin B6 Catalysis*. In: Korpela, T. and Christen, P. (Eds). Turku, Finland Birkhauser Verlag: 43-46.