

Immunofluorescence (Indirect Staining) Protocol for Adherent Cells

Ran Chen*

Department of Genetics, Stanford University, Stanford, USA

*For correspondence: rcchen@jfkbio.com

[Abstract] In this protocol, different types of adherent cells are fixed to coverslips by 4% paraformaldehyde. Target proteins are then stained by specific antibodies and fluorescent 2nd antibodies. This protocol therefore provides a method for immunostaining of adherent cells.

Materials and Reagents

1. Raw264.7, MCF-7 or HeLa cells
2. DPBS (Life Technologies, Invitrogen™, catalog number: 14190-250)
3. FBS (Atlanta Biologicals, catalog number: S11110H)
4. Anti-fade mounting medium: e.g. ProLong Gold Antifade Mountant (Life Technologies, Invitrogen™, catalog number: P10144) with DAPI (if nuclear staining is needed) or without DAPI
5. Paraformaldehyde (Sigma-Aldrich, catalog number: 158127)
6. General chemicals (Sigma-Aldrich)
7. Triton X-100 (Sigma-Aldrich, catalog number: T8787)
8. Tween 20 (Sigma-Aldrich, catalog number: P2287)
9. 4% paraformaldehyde-freshly-prepared (5 ml) (see Recipes)
10. Antibodies (see Recipes)

Equipment

1. Round cover slips (Thermo Fisher Scientific)
2. 24-well plate
3. Fluorescence microscope

Procedures

1. Grow cells on round cover slips in 24-well plates to 30-60% confluence. Do not overgrow because it will be difficult to distinguish cell components when stained.

2. Wash cells with 500 μ l/well DPBS twice, then add 250 μ l/well 4% paraformaldehyde for 15 min at room temperature (RT).
3. Remove paraformaldehyde, wash the fixed cells with 500 μ l/well PBS + 3% FBS for 3 times.
4. Permeabilize cells with 250 μ l/well DPBS + 0.2% Triton X-100 for 5 min, then wash with 500 μ l/well DPBS + 3% FBS for 3 times.
5. Block with 500 μ l/well DPBS + 3% FBS + 0.5% Tween 20 for 1 h at RT.
6. Remove the blocking buffer, add 250 μ l/well primary antibodies and incubate at RT for 1 h.
7. Remove the antibodies, wash with 500 μ l/well DPBS + 3% FBS in 5 min 3 times.
8. Add 250 μ l/well fluorescent secondary antibodies and incubate at RT for 30 min.
9. Remove the antibodies, wash with 500 μ l/well DPBS + 3% FBS in 5 min for 3 times.
10. Place a small drop of anti-fade reagent on a glass slide, then get the cover slip from the well and put it face down on the drop, push it tightly and attach to the glass slide.
11. Leave the slide in the dark for 5 min to let it dry.
12. The slide is ready to observe under a fluorescence microscope.

Recipes

1. 4% paraformaldehyde-freshly-prepared (5 ml)
 0.2 g paraformaldehyde powder + 5 ml DPBS + 50 μ l 1 N NaOH
 Incubate at 65 °C
 Vortex several times to dissolved completely
 Cool at room temperature then add 4 μ l HCl
 Mixed completely
2. Antibodies
 Diluted in DPBS + 3% FBS + 0.5 Tween 20
 Primary antibodies: 1:100-1:500 (depends on individual antibodies)
 Fluorescent secondary antibodies: 1:800

Acknowledgments

This work was funded by 5050 project by Hangzhou Hi-Tech District, Funding for Oversea Returnee by Hangzhou City, ZJ1000 project by Zhejiang Province. This protocol was developed in the Cohen Lab, Department of Genetics, Stanford University, CA, USA [Chen *et al.* (unpublished)].

References

1. Agrawal, S., van Dooren, G. G., Beatty, W. L. and Striepen, B. (2009). [Genetic evidence that an endosymbiont-derived endoplasmic reticulum-associated protein degradation \(ERAD\) system functions in import of apicoplast proteins.](#) *J Biol Chem* 284(48): 33683-33691.