

## Drug Sensitivity Assay of *Xanthomonas citri* subsp. *citri* Using REMA Plate Method

Isabel C. Silva\* and Henrique Ferreira

Ciências Biológicas, Universidade Estadual Paulista (UNESP), Araraquara, Brazil

\*For correspondence: [isabelcrs10@gmail.com](mailto:isabelcrs10@gmail.com)

**[Abstract]** Resazurin Microtiter Assay (REMA) is a simple, rapid, reliable, sensitive, safe and cost-effective measurement of cell viability. Resazurin detects cell viability by converting from a nonfluorescent dye to the highly red fluorescent dye resorufin in response to chemical reduction of growth medium resulting from cell growth (Palomino *et al.*, 2002). The REMA assay can be used as a fluorogenic oxidation-reduction indicator in a variety of cells, including bacteria, yeast and eukaryotes (Silva *et al.*, 2013).

### **Materials and Reagents**

1. Chemicals: Synthetic esters of gallic acids (Ximenes *et al.*, 2010)
2. Bacterial strain: Wild type *Xanthomonas citri* subsp *citri* strain 306 (Schaad *et al.*, 2005)
3. Dimethyl sulfoxide (DMSO) (Sigma-Aldrich, catalog number: D8418)
4. Kanamycin (Sigma-Aldrich, catalog number: K4000)
5. Luria-Bertani broth (LB) culture medium
6. Resazurin sodium salt (Sigma-Aldrich, catalog number: R7017)

### **Equipment**

1. 96-well plate, polystyrene, with clear flat bottom wells (Greiner Bio-one, catalog number: 655101)
2. SPECTRAfluor Plus (Tecan) microfluorimeter
3. Multichannel pipetman (Eppendorf)

### **Procedure**

- A. Prepare stock solutions of chemicals (dried-powder samples) dissolving in 10% in DMSO (diluted in sterile water).
- B. Add 100 µl of water to columns 1 and 12 to avoid evaporation (Table 1).


- C. Dilute the stock solutions in LB medium directly in a 96-well plates using a 2-fold scheme (final volume of 100  $\mu$ l per a well); after serial dilution, the most concentrated sample should have maximum 1% DMSO.
- D. Cells were grown in LB medium at 30 °C under rotation (200 rpm) until OD<sub>600</sub> 0.6 (log phase).
- E. Add 10  $\mu$ l of bacterial inoculum (standardized to 10<sup>5</sup> CFU/well).
  - a. Negative control: 1% DMSO dissolved in LB.
  - b. Positive control: Kanamycin at 15.6  $\mu$ g/ml.

**Table 1. Example for setup of REMA 96-well assay plate**

	1	2	3	4	5	6	7	8	9	10	11	12
A	200 $\mu$ l H <sub>2</sub> O	200 $\mu$ l drug 1	200 $\mu$ l drug 2	200 $\mu$ l drug 3	200 $\mu$ l drug 4	200 $\mu$ l drug 5	200 $\mu$ l drug 6	200 $\mu$ l drug 7	200 $\mu$ l drug 8	200 $\mu$ l drug 9	100 $\mu$ l negative control	200 $\mu$ l H <sub>2</sub> O
B	200 $\mu$ l H <sub>2</sub> O	100 $\mu$ l 2A	100 $\mu$ l 3A	100 $\mu$ l 4A	100 $\mu$ l 5A	100 $\mu$ l 6A	100 $\mu$ l 7A	100 $\mu$ l 8A	100 $\mu$ l 9A	100 $\mu$ l 10A	100 $\mu$ l negative control	200 $\mu$ l H <sub>2</sub> O
C	200 $\mu$ l H <sub>2</sub> O	100 $\mu$ l 2B	100 $\mu$ l 3B	100 $\mu$ l 4B	100 $\mu$ l 5B	100 $\mu$ l 6B	100 $\mu$ l 7B	100 $\mu$ l 8B	100 $\mu$ l 9B	100 $\mu$ l 10B	100 $\mu$ l negative control	200 $\mu$ l H <sub>2</sub> O
D	200 $\mu$ l H <sub>2</sub> O	100 $\mu$ l 2C	100 $\mu$ l 3C	100 $\mu$ l 4C	100 $\mu$ l 5C	100 $\mu$ l 6C	100 $\mu$ l 7C	100 $\mu$ l 8C	100 $\mu$ l 9C	100 $\mu$ l 10C	100 $\mu$ l negative control	200 $\mu$ l H <sub>2</sub> O
E	200 $\mu$ l H <sub>2</sub> O	100 $\mu$ l 2D	100 $\mu$ l 3D	100 $\mu$ l 4D	100 $\mu$ l 5D	100 $\mu$ l 6D	100 $\mu$ l 7D	100 $\mu$ l 8D	100 $\mu$ l 9D	100 $\mu$ l 10D	100 $\mu$ l positive control	200 $\mu$ l H <sub>2</sub> O
F	200 $\mu$ l H <sub>2</sub> O	100 $\mu$ l 2E	100 $\mu$ l 3E	100 $\mu$ l 4E	100 $\mu$ l 5E	100 $\mu$ l 6E	100 $\mu$ l 7E	100 $\mu$ l 8E	100 $\mu$ l 9E	100 $\mu$ l 10E	100 $\mu$ l positive control	200 $\mu$ l H <sub>2</sub> O
G	200 $\mu$ l H <sub>2</sub> O	100 $\mu$ l 2F	100 $\mu$ l 3F	100 $\mu$ l 4F	100 $\mu$ l 5F	100 $\mu$ l 6F	100 $\mu$ l 7F	100 $\mu$ l 8F	100 $\mu$ l 9F	100 $\mu$ l 10F	100 $\mu$ l positive control	200 $\mu$ l H <sub>2</sub> O
H	200 $\mu$ l H <sub>2</sub> O	100 $\mu$ l 2G	100 $\mu$ l 3G	100 $\mu$ l 4G	100 $\mu$ l 5G	100 $\mu$ l 6G	100 $\mu$ l 7G	100 $\mu$ l 8G	100 $\mu$ l 9G	100 $\mu$ l 10G	100 $\mu$ l positive control	200 $\mu$ l H <sub>2</sub> O

- F. Incubate the test plates at 30 °C for 6 h.
- G. Add 15  $\mu$ l of a 0.01% (w/v) resazurin solution, and incubate at 30 °C for 2 h.
- H. Measure fluorescence at 530 nm (excitation) and 590 nm (emission) using a fluorescence scanning.
- I. Percentage of inhibition is defined as:  

$$\frac{[(\text{average FU negative control}) - (\text{average FU test})]}{(\text{average FU negative control})} \times 100$$
 FU: Fluorescence Units



**Figure 1. Example for calculation of growth inhibition**

Note: Three independent experiments should be conducted, and the data is used to construct plots of chemical concentration versus cell growth inhibition in order to determine the MIC\* (Figure 1).

\*The minimum inhibitory concentration (MIC) is defined as the lowest concentration of the antibiotic able to inhibit the growth of 90% of organisms.

**Acknowledgments**

This work was supported by FAPESP research grants 2004/09173-6, 2010/05099-7, and 2011/07458-7. This protocol was adapted from a previous work by Palomino *et al.* (2002).

**References**

1. Palomino, J. C., Martin, A., Camacho, M., Guerra, H., Swings, J. and Portaels, F. (2002). [Resazurin microtiter assay plate: Simple and inexpensive method for detection of drug resistance in \*Mycobacterium tuberculosis\*](#). *Antimicrob Agents Chemother* 46(8): 2720-2722.
2. Schaad, N. W., Postnikova, E., Lacy, G. H., Sechler, A., Agarkova, I., Stromberg, P. E., Stromberg, V. K. and Vidaver, A. K. (2005). [Reclassification of \*Xanthomonas campestris\* pv. \*citri\* \(ex Hasse 1915\) Dye 1978 forms A, B/C/D, and E as \*X. smithii\* subsp. \*citri\* \(ex Hasse\) sp. nov. nom. rev. comb. nov., \*X. fuscans\* subsp. \*aurantifolii\* \(ex Gabriel 1989\) sp. nov. nom. rev. comb. nov., and \*X. alfalfae\* subsp. \*citrumelo\* \(ex Riker and Jones\) Gabriel \*et al.\*, 1989 sp. nov. nom. rev. comb. nov.; \*X. campestris\* pv. \*malvacearum\* \(ex smith 1901\) Dye 1978 as \*X. smithii\* subsp. \*smithii\* nov. comb. nov. nom. nov.; \*X. campestris\* pv.](#)

- [alfalfae \(ex Riker and Jones, 1935\) dye 1978 as \*X. alfalfae\* subsp. \*alfalfae\* \(ex Riker et al., 1935\) sp. nov. nom. rev.; and "var. fuscans" of \*X. campestris\* pv. \*phaseoli\* \(ex Smith, 1987\) Dye 1978 as \*X. fuscans\* subsp. \*fuscans\* sp. nov. Syst Appl Microbiol 28\(6\): 494-518.](#)
3. Silva, I. C., Regasini, L. O., Petronio, M. S., Silva, D. H., Bolzani, V. S., Belasque, J., Jr., Sacramento, L. V. and Ferreira, H. (2013). [Antibacterial activity of alkyl gallates against \*Xanthomonas citri\* subsp. \*citri\*](#). *J Bacteriol* 195(1): 85-94.
  4. Ximenes, V. F., Lopes, M. G., Petronio, M. S., Regasini, L. O., Silva, D. H. and da Fonseca, L. M. (2010). [Inhibitory effect of gallic acid and its esters on 2,2'-azobis\(2-amidinopropane\)hydrochloride \(AAPH\)-induced hemolysis and depletion of intracellular glutathione in erythrocytes](#). *J Agric Food Chem* 58(9): 5355-5362.